

Division Algebras: InterFamily Mixing (Including Neutrinos)

Geoffrey Dixon
gdixon@7stones.com

26 April 2003

It is shown how a 14-dimensional spacetime derived from the Division Algebras mixes leptoquark families due to the breaking of isospin $SU(2)$.

In [1] the ideas in [2,3] were extended. A one family leptoquark model founded on the Division Algebras was extended to a three family model by expanding the spinor space in which these fermions reside from $\mathbf{C} \otimes \mathbf{H}^2 \otimes \mathbf{O} = \mathbf{T}^2$ to

$$\mathbf{C} \otimes \mathbf{H}^2 \otimes \mathbf{O}^3 = \mathbf{T}^6,$$

which is acted upon by $\mathbf{T}_A(6)$ (see references for notation). $\mathbf{T}_A(6)$ is not a Clifford algebra, but contains three copies of $\mathcal{CL}(1, 13)$ (the Clifford algebra of 1,13-spacetime) which connect the families of the model in pairs.

Here we'll continue the ideas started in [1], with a focus on fermion interactions implied by a simple Dirac-like Lagrangian of the form

$$\mathcal{L} = \bar{\Psi} \not{\partial} \Psi,$$

where Ψ resides in \mathbf{T}^4 , and $\not{\partial}$ is the Dirac operator for $\mathcal{CL}(1, 13) \equiv \mathbf{T}_A(4)$.

First some notation: we use the 2×2 real matrices

$$\epsilon = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \alpha = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \beta = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \gamma = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Define, for example, the following 4×4 real matrix:

$$[\beta \otimes \alpha] = \begin{bmatrix} 0 & \alpha \\ \alpha & 0 \end{bmatrix}.$$

Our $\mathcal{CL}(1, 13)$ 1-vector basis consists of the following 14 elements:

1,3-spacetime:

$$[\epsilon \otimes \beta](iq_{R3}), [\epsilon \otimes \gamma]q_{Lk}e_{L7}(iq_{R3}), k = 1, 2, 3,$$

6 dimensional space; $su(3)$ charges; matter/antimatter mixing:

$$[\epsilon \otimes \gamma]ie_{Lp}(iq_{R3}), p = 1, \dots, 6,$$

4 dimensional space; $su(2)$ charges; interfamily mixing:

$$[\beta \otimes \epsilon]q_{R1}, [\beta \otimes \epsilon]q_{R2}, [\beta \otimes \alpha]q_{R3}, [\gamma \otimes \alpha].$$

It was shown in [3] how in a 1-family model based on $\mathcal{CL}(1, 9)$ a Dirac Lagrangian like that above gave rise to matter-antimatter mixing mediated by the extra 6 dimensions. (Note: particular fermions can be associated with particular bits of Ψ with the aid of projection operators; incoming fermions reside in Ψ , and outgoing in $\bar{\Psi}$; if a particular incoming/outgoing combination contributes a real part to \mathcal{L} , it is considered an observable interaction.) These extra dimensions also carried color $SU(3)$ charges, and by replacing the 6 real dimensions by 3 complex dimensions the offending terms of the Lagrangian disappeared (offending because we have not observed matter spontaneously transmuting into antimatter; disappeared because, for example, the complex partial $(\partial_x + i\partial_y)f(x + iy) = 0$ identically).

In the present case we have 4 dimensions beyond the original extra 6. These carry isospin $SU(2)$ charges and give rise to interfamily mixing. As we did in

the 6-dimensional case, if we replace the extra 4 dimensions with 2 complex dimensions we can make these mixing terms go away. However, $SU(3)$ is exact, and $SU(2)$ is broken, and it may not be justified to apply this method to all terms.

Begin by defining a pair of projection operators (idempotents) in $\mathbf{T}_A(4)$:

$$\Lambda_{\pm} = \frac{1}{2}(1 \pm iq_{R3}), \quad \Gamma_{\pm} = \frac{1}{2}(1 \pm [\alpha \otimes \epsilon]).$$

The two subspinors $\Gamma_{\pm}\Psi$ correspond to separate families (each is a 1,9-Dirac hyperspinor containing the 1,3-Dirac spinors of a family and its antifamily). The Λ_{\pm} are $SU(2)$ isospin projectors. The part of $\not{\partial}$ linear in the partials of these last 4 dimensions can be rewritten with respect to these projection operators as follows:

$$\begin{aligned} & ([\beta \otimes \epsilon]q_{R1}\partial_{10} + [\beta \otimes \epsilon]q_{R2}\partial_{11} + [\beta \otimes \alpha]q_{R3}\partial_{12} + [\gamma \otimes \alpha]\partial_{13}) \sum_{\pm\pm} (\Lambda_{\pm}\Gamma_{\pm}) \\ &= (q_{R1}[\beta \otimes \epsilon](\partial_{10} - i\partial_{11}) - i[\beta \otimes \alpha](\partial_{12} - i\partial_{13}))\Lambda_+\Gamma_+ \\ &+ (q_{R1}[\beta \otimes \epsilon](\partial_{10} + i\partial_{11}) + i[\beta \otimes \alpha](\partial_{12} + i\partial_{13}))\Lambda_-\Gamma_+ \\ &+ (q_{R1}[\beta \otimes \epsilon](\partial_{10} - i\partial_{11}) - i[\beta \otimes \alpha](\partial_{12} + i\partial_{13}))\Lambda_+\Gamma_- \\ &+ (q_{R1}[\beta \otimes \epsilon](\partial_{10} + i\partial_{11}) + i[\beta \otimes \alpha](\partial_{12} - i\partial_{13}))\Lambda_-\Gamma_- \end{aligned}$$

Meanwhile our Lagrangian term,

$$\mathcal{L} = \overline{\Psi} \not{\partial}\Psi = \begin{bmatrix} \Psi_{1r}^* & \Psi_{1l}^* & \Psi_{2r}^* & \Psi_{2l}^* \end{bmatrix} \not{\partial}_{1,13} \begin{bmatrix} \Psi_{1l} \\ \Psi_{1r} \\ \Psi_{2l} \\ \Psi_{2r} \end{bmatrix}$$

contains the following $\Lambda_+\Gamma_+$ part, linear in the partials of the extra 4 dimensions:

$$\mathcal{L}_{++} = \begin{bmatrix} \Psi_{1r}^* & \Psi_{1l}^* & \Psi_{2r}^* & \Psi_{2l}^* \end{bmatrix} \begin{bmatrix} (q_{R1}(\partial_{10} - i\partial_{11}) - i(\partial_{12} - i\partial_{13}))\Lambda_+\Psi_{2l} \\ (q_{R1}(\partial_{10} - i\partial_{11}) + i(\partial_{12} - i\partial_{13}))\Lambda_+\Psi_{2r} \\ 0 \\ 0 \end{bmatrix}$$

The subscripts 1 and 2 indicate different leptoquark families, while l and r the left and right Chiralities. The assumption we're making here is that as Ψ_{2l} is an $SU(2)$ doublet, its dependence on the extra 4 dimensions reduces to 2 complex dimensions in such a way that the partials in this Lagrangian term are identically zero. This leaves us with the term:

$$\Psi_{1l}^*(q_{R1}(\partial_{10} - i\partial_{11}) + i(\partial_{12} - i\partial_{13}))\Lambda_+\Psi_{2r}.$$

Since Ψ_{2r} is not an $SU(2)$ doublet we might assume that this term survives the partial, and since this is now associated with the term Ψ_{1l}^* , we infer that RH

members of family 2 mix via these extra dimensions with LH members of family 1. This would imply more than just neutrino oscillations: it would also imply interfamily mixing of charged leptons, and of quarks. It is unclear at present how probable any of these interactions might be. A single family/antifamily spinor is associated with 1,9-spacetime, 6 dimensions of which are tightly curled and linked to the exact internal symmetry, $U(1) \times SU(3)$. Adding a second family requires 4 more dimensions, also presumably curled, and linked to the broken symmetry, isospin $SU(2)$. (The third family makes things mathematically more complex [1].) A good many things are unclear at this point, but the structure is there to blow away much of the fog.

References:

- [1] G.M. Dixon, www.7stones.com/Homepage/6x6.pdf
- [2] G.M. Dixon, www.7stones.com/Homepage/10Dnew.pdf
- [3] G.M. Dixon, *Division Algebras: Octonions, Quaternions, Complex Numbers, and the Algebraic Design of Physics*, (Kluwer, 1994).